

GROUPS OF HIGHER DIMENSIONAL SATELLITE KNOTS

Taizo KANENOBU

*Department of Mathematics, Kobe University, Nada, Kobe, 657 Japan**

Communicated by H. Bass

Received June 1982

1. Introduction

The group of a knot is the fundamental group of the knot complement. A knot with a companion is called a satellite knot. For the classical knots, Schubert [21], [22] developed a theory of companionship of knots, which generalized the classical construction of composition of knots [20]. The higher dimensional satellite knots were first considered by Shinohara [25], who generalized the results on the Alexander polynomials and the signatures of classical satellite knots [23], [24] to the higher dimensional case. Recently, Yoshikawa [29] constructed an example of a non-fibered 2-knot which has a group with a nontrivial center using a satellite knot. In this paper, we investigate the groups of higher dimensional satellite knots.

In Section 2, the precise definition of a satellite knot is given, and the group of a satellite knot and its commutator subgroup are discussed. Section 3 contains an application of a satellite knot.

The deficiency of a 2-knot group is always ≤ 1 . Knots with deficiency one were constructed by Artin [2] and knots with deficiency zero by Fox [5]. Levine [15] have recently constructed knots of arbitrary large negative deficiency, by the composition of many knots of deficiency zero. In Section 3, we construct a satellite knot of any negative deficiency whose group is prime [17]. Modulo the unknotting conjecture for 2-knots, this is an affirmative answer to Problem 1.49 of Kirby [13], which was posed by S. Lomonaco.

The author is indebted to M. Sakuma for helpful conversations.

The remainder of this section is devoted to definitions and notational conventions.

We shall work throughout in the PL category. The boundary and interior of a manifold M are denoted by ∂M and $\text{int } M$. R^n, S^n and D^n denote, respectively, the Euclidean n -space, n -sphere and n -disk.

An n -knot K is a locally flat oriented submanifold of the oriented $(n+2)$ -sphere S^{n+2} which is homeomorphic to S^n . Throughout the paper we assume that the orientation of S^{n+2} is fixed.

Two n -knots K_1 and K_2 are *equivalent* if there exists an orientation preserving

* Currently at Department of Mathematics, Kyushu University, Fukuoka, 812 Japan.

homeomorphism f of S^{n+2} onto itself such that $f(K_1) = K_2$ and $f(K_1)$ is orientation preserving. An n -knot is *trivial* if it bounds an $(n+1)$ -disk in S^{n+2} .

Let G and H be groups. If G and H are isomorphic, then we write $G \cong H$. $G * H$ denotes the free product of G and H . The commutator subgroup of G is denoted by $[G]$.

Z denotes the infinite cyclic group or the integers. Z_n denotes the cyclic group of order n ; Z_0 means the infinite cyclic group Z . N denotes the natural numbers.

We write πK for the group of an n -knot K , i.e. $\pi K = \pi_1(S^{n+2} - K)$. $N(K)$ denotes a tubular neighborhood of K in S^{n+2} , which is homeomorphic to $S^n \times D^2$ [27, p. 268]. We call $x \in \pi K$ a *meridian* of K if x is represented by an oriented loop $p \times D^2$, $p \in S^n$, such that the intersection number of D^2 and K , regarding D^2 as a 2-chain and K as an n -cycle, is $+1$ in $H_0(S^{n+2}) \cong Z$.

Let K be an n -knot and $G = \pi K$. When we call G an n -knot group, we always consider the pair G and K , so we shall use the notation πK rather than G . Instead of this pair, we may consider the pair G and a meridian of K , see [17].

2. Satellite knots

Let J and K be n -knots and w and x be meridians of J and K , respectively. Any element of πK can be written in the form $x^p \beta$, where $\beta \in [\pi K]$, and can be represented by a simple closed curve l in $S^{n+2} - N(K)$ by Corollary 8.1.2 of [8]. Let V be a tubular neighborhood of l in $S^{n+2} - N(K)$. Since l bounds a 2-disk in S^{n+2} by Theorem 10.1 of [8], $S^{n+2} - \text{int } V$ is homeomorphic to $S^n \times D^2$. Let $h: S^{n+2} - \text{int } V \rightarrow N(J)$ be a homeomorphism. Then we obtain an n -knot $h(K)$. Note that $h(K)$ is not dependent on l , but on $x^p \beta$, because every simple closed curve representing $x^p \beta$ is contained in a unique ambient isotopy class by Theorem 10.1 of [8]. Therefore we denote $h(K)$ by $\Sigma(J; K, x^p \beta)$, and call it the *satellite knot of $(K, x^p \beta)$ about J* , or simply a *satellite knot*. We call J a *companion* of $\Sigma(J; K, x^p \beta)$. If either $x^p \beta = 1$ or J is trivial, then it is clear that $\Sigma(J; K, x^p \beta) = K$.

By the van Kampen Theorem, we have the following diagram of inclusion homomorphisms:

$$\begin{array}{ccc}
 & \pi_1(\partial N(J)) & \\
 i_1 \swarrow & & \searrow i_2 \\
 \pi_1(S^{n+2} - \text{int } N(J)) & & \pi_1(N(J) - h(K)) \\
 & \searrow & \swarrow \\
 & \pi_1(S^{n+2} - h(K)) &
 \end{array}$$

Clearly we have $\pi_1(\partial N(J)) = \langle w \rangle \cong Z$, $\pi_1(S^{n+2} - \text{int } N(J)) \cong \pi J$, $\pi_1(S^{n+2} - h(K)) = \pi \Sigma(J; K, x^p \beta)$ and $i_1(w) = w$. Furthermore, since an inclusion homomorphism

$$\pi_1((S^{n+2} - \text{int } V) - K) \rightarrow \pi K$$

is an isomorphism, we have

$$\pi_1(N(J) - h(K)) \cong \pi_1((S^{n+2} - \text{int } V) - K) \cong \pi K,$$

and $i_2(w) = x^p \beta$.

Let q be the order of $x^p \beta$ in πK ; if it is infinite, then let $q = 0$. Note that if $p \neq 0$ then $q = 0$, so $pq = 0$. Let $\langle w^q \rangle$ be the normal closure of w^q in πJ and $\pi J = \pi J / \langle w^q \rangle$. Then the order of w in $\tilde{\pi J}$ is q , since $\tilde{\pi J} / [\tilde{\pi J}] = \langle w \mid w^q = 1 \rangle \cong Z_q$. Then the subgroup of $\tilde{\pi J}$ generated by w , which we denote by H , and the subgroup of πK generated by $x^p \beta$ are both isomorphic to Z_q . Thus $\pi \Sigma(J; K, x^p \beta)$ is the free product of $\tilde{\pi J}$ and πK with an amalgamated subgroup H [18, p. 207]:

$$\pi \Sigma(J; K, x^p \beta) = \tilde{\pi J} *_H \pi K,$$

and so it contains subgroups isomorphic to $\tilde{\pi J}$ and πK .

Now we write down a presentation of $\pi \Sigma(J; K, x^p \beta)$ in terms of presentations of $[\pi J]$, $[\pi K]$, πJ and πK . Let

$$[\pi J] = \langle A \mid T \rangle \quad \text{and} \quad [\pi K] = \langle B \mid U \rangle,$$

where

$$\begin{aligned} A &= \{a_\mu \mid \mu \in N\}, & B &= \{b_\nu \mid \nu \in N\}, \\ T &= \{t_i(a_\mu) \mid i \in N\}, & U &= \{u_j(b_\nu) \mid j \in N\}; \end{aligned}$$

$t_i(a_\mu)$ is a word in A and $u_j(b_\nu)$ a word in B . Since πJ is a split extension of $[\pi J]$ by the infinite cyclic group $\pi J / [\pi J] = \langle w \mid \rangle$, πJ has the following presentation [7, 15.4]:

$$\pi J = \langle w, A \mid R, T \rangle, \quad *$$

where

$$R = \{w a_i w^{-1} = r_i(a_\mu) \mid i \in N\};$$

$r_i(a_\mu)$ is a word in A . Similarly, we have

$$\pi K = \langle x, B \mid S, U \rangle,$$

where

$$S = \{x b_j x^{-1} = s_j(b_\nu) \mid j \in N\};$$

$s_j(b_\nu)$ is a word in B . Hence

$$\pi \Sigma(J; K, x^p \beta) = \langle w, x, A, B \mid R, S, T, U, w = x^p \beta \rangle.$$

Theorem 2.1. *If $p > 0$, then*

$$[\pi \Sigma(J; K, x^p \beta)] = \bigast_{k=0}^{p-1} x^k [\pi J] x^{-k} * [\pi K],$$

and if $p < 0$, then

$$[\pi \Sigma(J; K, x^p \beta)] = \bigast_{k=p+1}^0 x^k [\pi J] x^{-k} * [\pi K].$$

Proof. Suppose that $p > 0$. Let $\alpha_{\mu_k} = x^k a_{\mu} x^{-k}$ and

$$A_k = \{\alpha_{\mu_k} \mid \mu \in N\}.$$

Since

$$\begin{aligned} w a_i w^{-1} &= (x^p \beta x^{-p}) \{x(x^{p-1} a_i x^{-(p-1)}) x^{-1}\} (x^p \beta^{-1} x^{-p}) \\ &= (x^p \beta x^{-p}) (x \alpha_{i, p-1} x^{-1}) (x^p \beta^{-1} x^{-p}) \end{aligned}$$

and

$$a_{\mu} = \alpha_{\mu_0},$$

we have

$$x \alpha_{i, p-1} x^{-1} = (x^p \beta^{-1} x^{-p}) r_i(\alpha_{\mu_0}) (x^p \beta x^{-p}).$$

Let

$$\begin{aligned} \tilde{R} = \{x \alpha_{ik} x^{-1} = \alpha_{i, k+1}, x \alpha_{i, p-1} x^{-1} = (x^p \beta^{-1} x^{-p}) r_i(\alpha_{\mu_0}) (x^p \beta x^{-p}) \mid \\ i \in N, k = 0, 1, \dots, p-2\}. \end{aligned}$$

Let

$$T_k = \{t_i(\alpha_{\mu_k}) \mid i \in N\},$$

where $t_i(\alpha_{\mu_k}) = x^k t_i(a_{\mu}) x^{-k}$. Then we have

$$\pi \Sigma(J; K, x^p \beta) = \langle x, A_0, \dots, A_{p-1}, B \mid \tilde{R}, S, T_0, \dots, T_{p-1}, U \rangle.$$

Thus we obtain

$$\begin{aligned} [\pi \Sigma(J; K, x^p \beta)] &= \langle A_0, \dots, A_{p-1}, B \mid T_0, \dots, T_{p-1}, U \rangle \\ &= \langle A_0 \mid T_0 \rangle * \dots * \langle A_{p-1} \mid T_{p-1} \rangle * \langle B \mid U \rangle. \end{aligned}$$

Since $\langle A_k \mid T_k \rangle = x^k [\pi J] x^{-k}$, the result follows. For the case $p < 0$, we can show in the same way. \square

If L is a fibered n -knot, then $[\pi L]$ is finitely generated [19, p. 324]. Yoshikawa [29] proved:

Proposition 2.2. *If $\tilde{\pi} J \not\cong Z_q$, $\pi K \not\cong Z$ and $\beta \neq 1$, then $[\pi \Sigma(J; K, \beta)]$ is not finitely generated.*

In contrast with this proposition, we show the following theorem, which is a generalization of Theorem 5 of [26] to higher dimensions.

Theorem 2.3. *If J and K are fibered knots with fibers E and F , respectively and $p \neq 0$, then $\Sigma(J; K, x^p \beta)$ is a fibered knot with fiber a connected sum of $|p|$ copies of E and F ; $\#_p E \# F$.*

Proof. Let l be a simple closed curve in $S^{n+2} - N(K)$ representing $x^p \beta$. We may suppose that l meets F transversely in $|p|$ points. Then $(S^{n+2} - \text{int } V) - K$ fibers over S^1 with fiber $F - \text{int } V$, where V is a tubular neighborhood of l in $S^{n+2} - N(K)$. Thus the fiber of $\Sigma(J; K, x^p \beta)$ can be constructed by piecing together $|p|$ copies of E with $F - \text{int } V$. \square

A satellite knot $\Sigma(J; K, x)$ is the *composition of J and K* and denoted by $J \# K$, see [27, pp. 342–343]. An n -knot L is *prime* if $L = J \# K$ implies that either J or K is a trivial n -knot. We call $\pi(J \# K)$ the *composition of πJ and πK* , and denote it by $\pi J \# \pi K$ [17, p. 591]. An n -knot group πL is *prime* if $\pi L = \pi J \# \pi K$ implies either πJ or πK is isomorphic to Z . For $n \geq 3$, an n -knot L is not always prime though πL is prime, because there exists a nontrivial n -knot L with $\pi L \cong Z$ [14], [19, p. 285]. For 2-knots, modulo the unknotting conjecture [27, p. 342], a 2-knot L is prime if πL is prime.

By Theorem 2.1, we have immediately the following proposition due to Maeda [17, p. 592].

Proposition 2.4. *Let $\pi L = \pi J \# \pi K$, and let $\langle [\pi J] \rangle$ and $\langle [\pi K] \rangle$ be the normal closures of $[\pi J]$ and $[\pi K]$ in πL , respectively. Then*

- (1) $[\pi L] = [\pi J] * [\pi K]$;
- (2) $\pi J = \pi L / \langle [\pi K] \rangle$ and $\pi K = \pi L / \langle [\pi J] \rangle$.

3. The deficiency of a 2-knot groups

The deficiency of a finite presentation

$$\langle x_1, \dots, x_p \mid r_1, \dots, r_q \rangle$$

of a group G is defined as $p - q$. The *deficiency of a group G* , denoted by $\text{def } G$, is the maximum of the deficiencies of its finite presentations.

Kervaire [11], [12] showed that any knot group G has the following properties:

- (i) G is finitely presented.
- (ii) $G/[G] \cong Z$.
- (iii) The normal closure of some single element is G .
- (iv) $H_2(G; Z) = 0$.

Conversely, he showed that for any $n \geq 3$, any group G with properties (i)–(iv) is the group of some n -knot. For $n = 2$, he proved that if G satisfies (i)–(iii) and the additional property:

- (iv)' $\text{def } G = 1$,

then there exists a homotopy 4-sphere Σ^4 and a locally flat embedding $K : S^2 \rightarrow \Sigma^4$ such that $G = \pi_1(\Sigma^4 - K(S^2))$.

If G is a knot group, then $\text{def } G \leq 1$ [4]. Every spun 2-knot [2], [1], [19, p. 85] has a group with deficiency one, since its group is isomorphic to a 1-knot group. The 2-knot of Example 12 in [5] has a group with deficiency zero [11, p. 106]. Levine [15] constructed 2-knots of deficiency < 0 ; he showed that $\text{def } \pi F_m = 1 - m$, where F_m is the composition of m copies of the 2-twist spun trefoil [30].

Remark. The 2-knot of Example 12 in [5] and the 2-twist spun trefoil are equivalent [16], [10].

Accordingly Lomonaco asked in [13, Problem 1.49] whether a prime 2-knot can have deficiency < 0 . If the unknotting conjecture for 2-knots is correct, then Theorem 3.2 below gives an affirmative answer of this problem.

Lemma 3.1. (Compare [28, p. 322].) *Let T be the 3-twist spun trefoil. Let*

$$P_1 = \langle x, a \mid xax = ax^2a, x^3 = (ax)^3 \rangle$$

and

$$P_2 = \langle x, a, b \mid xax^{-1} = b, xbx^{-1} = ba, a^2 = b^2 = (ab)^2 \rangle.$$

Then $\pi T \cong P_1 \cong P_2$, where x is a meridian, and

$$\{\pi T\} = \langle a, b \mid a^2 = b^2 = (ab)^2 \rangle,$$

which is the quaternion group of order 8.

Proof. By [30], we have

$$\pi T = \langle x, y \mid xyx = yxy, y = x^{-3}yx^3 \rangle,$$

where x and y are meridians. This presentation indicates that x^3 is in the center of πT . From the first relation, we have $yx y^{-1} = x^{-1}yx$, and so $yx^3 y^{-1} = x^{-1}y^3x$. Thus we have $x^3 = y^3$. Therefore

$$\pi T = \langle x, y \mid xyx = yxy, x^3 = y^3 \rangle.$$

Letting $y = ax$, we obtain $\pi T \cong P_1$.

Let $b = xax^{-1}$. From the second relation of P_1 , we have $a(xax^{-1})(x^2ax^{-2}) = 1$, and so $xbx^{-1} = x^2ax^{-2} = b^{-1}a^{-1}$. From the first relation of P_1 , we have $b = a(b^{-1}a^{-1})$ and $b^{-1}a^{-1} = ba$. Hence we obtain $\pi T \cong P_2$. \square

Remark. πT is isomorphic to the group of the roll-spun figure eight knot [6]. Also πT is isomorphic to the group of K_1 in [9]. The question of whether these three knots are equivalent is open.

Theorem 3.2. *Let T be the 3-twist spun trefoil and*

$$\pi T = \langle x, a \mid xax = ax^2a, x^3 = (ax)^3 \rangle.$$

Let T_m be the composition of m copies of the 3-twist spun trefoil. Let $K_m = \Sigma(T_m; T, x^2a)$. Then

- (1) $\text{def } \pi K_m = -m$;
- (2) πK_m is prime.

Proof. (1) πT_m has a presentation

$$\langle w, c_1, \dots, c_m \mid wc_iw = c_iw^2c_i, w^3 = (c_iw)^3, i = 1, \dots, m \rangle.$$

Then πK_m is presented by

$$G_1 = \langle x, a, c_1, \dots, c_m \mid xax = ax^2a, x^3 = (ax)^3, x^2ac_ix^2a = c_i(x^2a)^2c_i, \\ (x^2a)^3 = (c_ix^2a)^3, i = 1, \dots, m \rangle.$$

Thus $\text{def } \pi K_m \geq -m$.

Let \tilde{X}_m be the infinite cyclic covering space of $S^4 - K_m$. Then $H_*(\tilde{X}_m)$ has a Λ -module structure, where Λ is the polynomial ring $Z[t, t^{-1}]$ [19, Chapter 7]. In [15], $H_*(\tilde{X}_m)$ is denoted by $H_*(\pi K_m; \Lambda)$. From the above presentation of πK_m , the method of [19, Chapter 7] shows that

$$H_1(\tilde{X}_m) = A \oplus \underbrace{B \oplus \dots \oplus B}_m,$$

where $A = \Lambda/(2, t^2 + t + 1)$ and $B = \Lambda(2, t^4 + t^2 + 1)$. If we regard A and B as Λ_2 -modules, where $\Lambda_2 = \Lambda/(2)$, then $A = \Lambda_2/(t^2 + t + 1)$ and $B = \Lambda_2/(t^4 + t^2 + 1)$. Note that $t^4 + t^2 + 1 = (t^2 + t + 1)(t^2 - t + 1)$. By the structure theorem for modules over the principal ideal domain Λ_2 , we can see that $H_1(\tilde{X}_m)$ cannot be generated as a Λ -module by fewer than $p + 1$ elements. Applying the proof of Theorem II of [15], we can show that $\text{def } \pi K_m \leq -m$, so the result holds.

(2) By Lemma 3.1, letting $b = xax^{-1}$ and $d_i = xc_ix^{-1}$, we have

$$\pi T = \langle x, a, b \mid xax^{-1} = b, xbx^{-1} = ba, a^2 = b^2 = (ab)^2 \rangle$$

and

$$\pi T_m = \langle w, c_1, d_1, \dots, c_m, d_m \mid wc_iw^{-1} = d_i, wd_iw^{-1} = d_ic_i, \\ c_i^2 = d_i^2 = (c_id_i)^2, i = 1, \dots, m \rangle.$$

Thus by Theorem 2.1, we have

$$[\pi K_m] = [\pi T_m] * x[\pi T_m]x^{-1} * [\pi T],$$

where

$$[\pi T] = \langle a, b \mid a^2 = b^2 = (ab)^2 \rangle$$

and

$$[\pi T_m] = \langle c_1, d_1, \dots, c_m, d_m \mid c_i^2 = d_i^2 = (c_id_i)^2, i = 1, \dots, m \rangle.$$

Let $Q_0 = [\pi T]$, $Q_{2i-1} = \langle c_i, d_i \mid c_i^2 = d_i^2 = (c_id_i)^2 \rangle$ and $Q_{2i} = xQ_{2i-1}x^{-1}$. Then

$$[\pi K_m] = \bigstar_{j=0}^{2m} Q_j.$$

Here each Q_j is isomorphic to the quaternion group, so Q_j is indecomposable relative to free products [18, p. 195].

Now we prove the primeness of πK_m . Assume that πK_m is not prime; $\pi K_m = \pi J_1 \# \pi J_2$, where neither πJ_1 nor πJ_2 is infinite cyclic group. Then by Proposition 2.4, $[\pi K_m] = [\pi J_1] * [\pi J_2]$. Since Q_{2i-1} and Q_{2i} are conjugate in πK_m , we may suppose that

$$[\pi J_1] = \bigstar_{j=0}^{2l-2} Q_j \quad \text{and} \quad [\pi J_2] = \bigstar_{j=2l-1}^{2m} Q_j,$$

where $1 \leq l \leq m$. By Proposition 2.4, we have

$$\begin{aligned} \pi J_1 = \langle x, a, c_1, \dots, c_{l-1} \mid xax = ax^2a, x^3 = (ax)^3, x^2ac_ix^2a = c_i(x^2a)^2c_i, \\ (x^2a)^3 = (c_ix^2a)^3, i = 1, \dots, l-1 \rangle \end{aligned}$$

and

$$\pi J_2 = \langle x, c_l, \dots, c_m \mid x^2c_jx^2 = c_jx^4c_j, x^6 = (c_jx^2)^3, j = l, \dots, m \rangle$$

Thus $\pi K_m = \pi J_1 \# \pi J_2$ is presented by

$$\begin{aligned} G_2 = \langle x, a, c_1, \dots, c_m \mid xax = ax^2a, x^3 = (ax)^3, \\ x^2ac_ix^2a = c_i(x^2a)^2c_i, (x^2a)^3 = (c_ix^2a)^3, \\ x^2c_jx^2 = c_jx^4c_j, x^6 = (c_jx^2)^3, \\ i = 1, \dots, l-1, j = l, \dots, m \rangle. \end{aligned}$$

Therefore G_1 and G_2 must be isomorphic under the mapping $x \rightarrow x, a \rightarrow a, c_i \rightarrow c_i$.

Adding to G_1 and G_2 the relations $x^3 = c_1 = \dots = c_{m-1}$, and letting $c = c_m$, we obtain

$$\tilde{G}_1 = \langle x, a, c \mid xax = ax^2a, x^3 = (ax)^3 = 1, x^2acx^2a = c(x^2a)^2c, (x^2a)^3 = (cx^2a)^3 \rangle$$

and

$$\tilde{G}_2 = \langle x, a, c \mid xax = ax^2a, x^3 = (ax)^3 = 1, x^2cx^2 = cx^4c, x^6 = (cx^2)^3 \rangle,$$

respectively. These two groups must be isomorphic under the mapping $x \rightarrow x, a \rightarrow a, c \rightarrow c$.

First, we consider \tilde{G}_1 . Letting $y = ax$ and $z = x^2a$, and eliminating a , we obtain

$$\tilde{G}_1 = \langle x, y, z, c \mid xyx = yxy, x^3 = y^3 = 1, zcx = cz^2c, z^3 = (cz)^3, z = x^2yx^2 \rangle.$$

Let

$$B_1 = \langle x, y \mid xyx = yxy, x^3 = y^3 = 1 \rangle.$$

Then by Lemma 3.1, B_1 is a factor group of the group of the 3-twist spun trefoil, and is isomorphic to

$$\langle x, y \mid xyx = yxy, x^3 = 1 \rangle,$$

which is the binary tetrahedral group of order 24 [3, p. 134], and is also isomorphic to the special linear group $SL(2, 3)$, the group of all 2×2 matrices of integers (mod 3) of determinant 1, under the mapping $x \rightarrow \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ and $y \rightarrow \begin{pmatrix} 1 & 0 \\ 2 & 1 \end{pmatrix}$, see [3, p. 98]. Then x^2yx^2 is mapped to $\begin{pmatrix} 2 & 0 \\ 2 & 2 \end{pmatrix}$, so it has order 6 in B_1 . Let

$$A_1 = \langle z, c \mid zcz = cz^2c, z^3 = (cz)^3, z^6 = 1 \rangle.$$

Then $A_1/[A_1] = \langle z \mid z^6 = 1 \rangle \cong Z_6$. Thus \tilde{G}_1 is the free product of A_1 and B_1 with an amalgamated subgroup $H_1 = \langle z \mid z^6 = 1 \rangle \cong Z_6$:

$$\tilde{G}_1 = A_1 *_H B_1.$$

Let us denote the center of a group G by $C(G)$. By Corollary 4.5 of [18], $C(\tilde{G}_1) = H_1 \cap C(A_1) \cap C(B_1)$. Since the center of $SL(2, 3)$ is

$$\left\{ \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 2 & 0 \\ 0 & 2 \end{pmatrix} \right\} \cong Z_2$$

and $(x^2yx^2)^3 (=z^3)$ is mapped to $\begin{pmatrix} 2 & 0 \\ 0 & 2 \end{pmatrix}$, $C(\tilde{G}_1) \cong Z_2$.

Next, we consider \tilde{G}_2 . We have

$$\tilde{G}_2 = \langle x, a, c \mid xax = ax^2a, x^2cx^2 = cxc, x^3 = (ax)^3 = (cx^2)^3 = 1 \rangle.$$

Let

$$A_2 = \langle x, c \mid x^2cx^2 = cxc, x^3 = (cx^2)^3 = 1 \rangle,$$

$$B_2 = \langle x, a \mid xax = ax^2a, x^3 = (ax)^3 = 1 \rangle.$$

Then it is easy to see that both A_2 and B_2 are isomorphic to the binary tetrahedral group. Hence \tilde{G}_2 is the free product of A_2 and B_2 with an amalgamated subgroup $H_2 = \langle x \mid x^3 = 1 \rangle \cong Z_3$:

$$\tilde{G}_2 = A_2 *_{{H_2}} B_2.$$

Thus as before $C(\tilde{G}_2) = H_2 \cap C(A_2) \cap C(B_2)$, which is a trivial group, because $H_2 \cong Z_3$ and $C(A_2) \cong C(B_2) \cong Z_2$.

Hence \tilde{G}_1 and \tilde{G}_2 are not isomorphic, and this contradiction establishes the primeness of πK_m . \square

References

- [1] J.J. Andrews and M.L. Curtis, Knotted 2-spheres in the 4-sphere, *Ann. of Math.* 70 (1959) 565–571.
- [2] E. Artin, Zur Isotopie zweidimensionaler Flächen im R_4 , *Abh. Math. Sem. Univ. Hamburg* 4 (1925) 174–177.
- [3] H.S.M. Coxeter and W.O.J. Moser, *Generators and Relations for Discrete Groups* (Springer, Berlin-Heidelberg-New York, 1980).
- [4] D.B.A. Epstein, Finite presentations of groups and 3-manifolds, *Quart. J. Math.* 12 (1961) 205–212.
- [5] R.H. Fox, A quick trip through knot theory, in: *Topology of 3-Manifolds and Related Topics* (Prentice-Hall, Englewood Cliffs, 1962) 120–167.
- [6] R.H. Fox, Rolling, *Bull. Amer. Math. Soc.* 72 (1966) 162–164.
- [7] M. Hall, Jr., *The Theory of Groups* (Macmillan, New York, 1959).
- [8] J.F.P. Hudson, *Piecewise Linear Topology* (Benjamin, New York, 1969).
- [9] T. Kanenobu, 2-knot groups with elements of finite order, *Math. Sem. Notes Kobe Univ.* 8 (1980) 557–560.
- [10] T. Kanenobu, Fox's 2-spheres are twist spun knots, Preprint (1981).
- [11] M.A. Kervaire, On higher dimensional knots, in: *Differential and Combinatorial Topology* (Princeton Univ. Press, Princeton, 1965) 105–120.
- [12] M.A. Kervaire, Les noeuds de dimension supérieures, *Bull. Soc. Math. France* 93 (1965) 225–271.
- [13] R. Kirby, Problems in low-dimensional manifold theory, in: *Proc. Symp. Pure Math. XXXII, Part 2* (Amer. Math. Soc., Providence, RI, 1978) 273–312.
- [14] J. Levine, An algebraic classification of some knots of codimension two, *Comment. Math. Helv.* 45 (1970) 185–198.

- [15] J. Levine, Some results on higher dimensional knot theory, in: *Knot theory*, Lecture Notes in Mathematics No. 685 (Springer, Berlin-Heidelberg-New York, 1978) 243-269.
- [16] R.A. Litherland, *Topics in Knot Theory*, Ph.D. Thesis, Cambridge University (1978).
- [17] T. Maeda, A unique decomposition for knot-like groups, *Math. Sem. Notes Kobe Univ.* 6 (1978) 567-602.
- [18] W. Magnus, A. Karrass and D. Solitar, *Combinatorial Group Theory* (Dover, New York, 1976).
- [19] D. Rolfsen, *Knots and Links*, Publish or Perish Math. Lecture Series 7 (Publish or Perish, Berkeley, CA, 1976).
- [20] H. Schubert, Die eindeutige Zerlegbarkeit eines Knotens in Primknoten, *S.-B. Heidelberger Akad. Wiss. Math.-Nat. Kl.* 3 (1949) 57 -104.
- [21] H. Schubert, Knoten und Vollringe, *Acta. Math.* 90 (1953) 131-286.
- [22] H. Schubert, Über eine numerische Knoteninvariante, *Math. Z.* 61 (1954) 245-288.
- [23] H. Seifert, On the homology invariants of knots, *Quart. J. Math.* 1 (1950) 23-32.
- [24] Y. Shinohara, On the signature of knots and links, *Trans. Amer. Math. Soc.* 156 (1971) 273-285.
- [25] Y. Shinohara, Higher dimensional knots in tubes, *Trans. Amer. Math. Soc.* 161 (1971) 35-49.
- [26] J. Stallings, Constructions of fibred knots and links, in: *Proc. Symp. Pure Math. XXXII*, Part 2 (Amer. Math. Soc., Providence RI, 1978) 55-60.
- [27] S. Suzuki, Knotting problems of 2-spheres in 4-sphere, *Math. Sem. Notes Kobe Univ.* 4 (1976) 241-371.
- [28] K. Yoshikawa, On 2-knot groups with the finite commutator subgroups, *Math. Sem. Notes Kobe Univ.* 8 (1980) 321-330.
- [29] K. Yoshikawa, On a 2-knot group with nontrivial center, *Bull. Austral. Math. Soc.* 25 (1982) 321-326.
- [30] E.C. Zeeman, Twisting spun knots, *Trans. Amer. Math. Soc.* 115 (1965) 471-495.